


CDC Plentiful

Wheat – CWRS

CDC Plentiful offers high yield, strong straw and a comprehensive disease package including the best available resistance (MR) to fusarium head blight. In addition, this variety is very early maturing (1 to 3 days earlier than most popular varieties e.g. Cardale, Carberry), making it ideal for all regions.



YIELD PERFORMANCE



Source: seed.ab.ca, Sask. Seed 2015, Seed Manitoba 2015

Parentage: BW282/CDC Go

Yield
(Western Canada)

106 – 109%*

Maturity

- 3 days earlier than check

Disease Resistance

- MR rating to FHB
- R to leaf and stem rust, MR to stripe rust
- I to bunt

Harvestability

- Easy to thresh
- Strong straw

* Percentage of check



Higher yield and more profit guaranteed for this variety! Ask us for details.



CDC Plentiful

Wheat – CWRS

2015 ALBERTA SEED GUIDE AGRONOMIC DATA

| Variety | Yield (% of AC Barrie) | Protein (+/- check) | Maturity | Lodging | Height (cm) | Awned | Resistance to | | |
|----------------------|---------------------------|------------------------|----------|-----------|----------------|-----------|---------------|-------------|-----------|
| | | | | | | | Sprouting | Stripe Rust | FHB |
| AC Barrie | 59 bu/ac. | 14.1 | M | G | 89 | No | G | S | I |
| AC® Harvest | 102 | -0.1 | M | VG | 84 | No | VG | MR | S |
| Lillian | 104 | 0.2 | M | F | 86 | No | G | R | S |
| AC® Muchmore | 111 | -0.9 | L | VG | 75 | Yes | G | MR | MS |
| Carberry | 108 | -0.6 | L | VG | 79 | Yes | F | MR | MR |
| Stettler | 112 | -0.3 | M | G | 84 | Yes | G | I | MS |
| CDC Go | 110 | -0.1 | M | G | 83 | Yes | VP | MR | MS |
| CDC Utmost VB | 112 | -0.2 | M | G | 85 | No | G | I | MS |
| CDC Plentiful | 106 | -0.2 | M | VG | 87 | No | P | MR | MR |

Check is AC Barrie: average yield = 59 bu/ac; protein = 14.1 %

Source: seed.ab.ca

2015 SASKATCHEWAN SEED GUIDE AGRONOMIC DATA

| Variety | Yield (% of Carberry) | Protein (+/- check) | Maturity (+/- Carberry) | Lodging | Height (cm) (+/- check) | Awned | Resistance to | | |
|----------------------|--------------------------|------------------------|----------------------------|-----------|----------------------------|-----------|---------------|----------|-----------|
| | | | | | | | Sprouting | Bunt | FHB |
| Carberry | 100 | 14.6 | 105 days | VG | 84 | Yes | I | R | MR |
| AC® Harvest | 99 | -0.3 | -3 | VG | 10 | No | R | S | S |
| Lillian | 96 | 1.1 | -2 | F | 12 | No | MR | MR | S |
| AC® Muchmore | 101 | -0.4 | 0 | VG | -3 | Yes | MR | R | MS |
| AC Barrie | 97 | 0.1 | -2 | G | 12 | No | MR | I | I |
| Stettler | 106 | 0.2 | -1 | G | 7 | Yes | MR | MR | MS |
| CDC Utmost VB | 110 | -0.4 | -3 | G | 10 | No | MR | S | MS |
| CDC Plentiful | 107 | -0.3 | -3 | VG | 8 | No | MS | I | MR |

Source: Sask Seed 2015

2015 MANITOBA SEED GUIDE AGRONOMIC DATA

| Variety | Yield (bu/ac) | Protein (%) | Maturity (+/- 99 days) | Lodging | Height (+/- 37 in) | Awned | Resistance to | | |
|----------------------|------------------|----------------|---------------------------|-----------|-----------------------|-----------|---------------|----------|-----------|
| | | | | | | | Leaf Rust | Bunt | FHB |
| AC Barrie | 57 | 14.6 | 0 | G | 0 | No | MS | I | I |
| AC® Harvest | 60 | 14.8 | -1 | VG | -2 | No | MR | S | S |
| AC® Muchmore | 62 | 14.4 | 2 | VG | -6 | Yes | R | R | MS |
| Carberry | 62 | 14.5 | 2 | VG | -5 | Yes | R | R | MR |
| Cardale | 64 | 14.6 | 2 | VG | -4 | Yes | R | MR | MR |
| Kane | 61 | 14.4 | 1 | G | -2 | Yes | R | I | I |
| Glenn | 62 | 14.1 | 2 | VG | -2 | Yes | R | I | I |
| Stettler | 59 | 14.8 | 1 | G | -2 | Yes | MS | MR | MS |
| CDC Utmost VB | 63 | 14.4 | -1 | G | -1 | No | R | S | MS |
| CDC Plentiful | 62 | 14.1 | -3 | VG | -2 | No | R | I | MR |

Source: Seed Manitoba 2015